

**COVID-19 VACCINE IMMUNIZATION CONSENT FORM**

**For COVID-19 Provider use only** Clinic Name/Code: Southern Pharmacy of Arkansas  
 Location type: Pharmacy  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ County: \_\_\_\_\_  
 State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

**Person Receiving Vaccine:**

**(Legal) First Name:** \_\_\_\_\_ **MI:** \_\_\_\_ **Last Name:** \_\_\_\_\_

**Date of Birth:**   /   /

**MEDICAL HISTORY: Complete the following questions for the individual receiving the vaccine.**

**If you answer "YES" you may not be able to receive the COVID-19 vaccine.**

<i>If YES refer to Pfizer website at <a href="http://www.PfizerMedInfo.com">www.PfizerMedInfo.com</a>. For Moderna COVID-19 vaccine <a href="http://www.modernatx.com">www.modernatx.com</a>. For Janssen COVID-19 vaccine <a href="http://www.janssencovid19vaccine.com">www.janssencovid19vaccine.com</a>. Refer to Pre-vaccination Checklist for COVID-19 Vaccines Information for Healthcare Professionals (<a href="http://cdc.gov">cdc.gov</a>) to clarify further questions. <a href="http://www.cdc.gov/vaccines/covid-19/downloads/pre-vaccination-screening-form.pdf">www.cdc.gov/vaccines/covid-19/downloads/pre-vaccination-screening-form.pdf</a>.</i>	<b>YES</b>	<b>NO</b>
Have you had a previous COVID-19 vaccine? If yes, date?		
Have you had any vaccines within the previous 14 days? Pfizer-BioNTech, Moderna, or Janssen COVID-19 vaccine should be administered alone with minimal interval of 14 days before or after any other vaccine.		
Do you have a fever today? Are you sick today? Do you have COVID-19 infection and are currently in isolation? Are you currently in quarantine for known exposure to COVID-19?		
Have you ever had severe allergic reaction (anaphylactic reaction that required treatment with EpiPen or treatment at a hospital) to any vaccine (including Pfizer-BioNTech, Moderna, or Janssen COVID-19 vaccine), or vaccine component (including polyethylene glycol [PEG] in medications or laxatives and preparations for colonoscopy), or immediate allergic reaction of any severity to polysorbate in vaccines, coated tablets or IV steroids (due to potential cross-reactive hypersensitivity with the vaccine ingredient PEG) or injectable therapy? This would include an allergic reaction that occurred within 4 hours, such as difficulty breathing, hives, swelling of your face and throat, fast heartbeat, bad rash all over your body, dizziness, and weakness.		
Have you ever had a severe allergic reaction (anaphylaxis) to something other than a component of COVID-19 vaccine or any vaccine or injectable medication such as food, pet, venom, environmental, or oral medication allergies?		
Do you have a bleeding disorder or are you taking a blood thinner? If so, a fine gauge needle (23 gauge or smaller caliber) should be used to administer the vaccine, followed by firm pressure without rubbing for at least 2 minutes.		
Do you have dermal fillers? If swelling occurs at or near the filler injection site, usually face or lips, patient should contact their health care provider.		
Are you pregnant, breastfeeding, or planning to become pregnant? Women in this group may receive Pfizer-BioNTech, Moderna, or Janssen COVID-19 vaccine. A discussion with your doctor can help make an informed decision		
Are you immunocompromised? Do you have a condition that weakens your immune system? Are you receiving any immunosuppressive therapy? You are still eligible to receive Pfizer-BioNTech, Moderna, or Janssen COVID-19 vaccine unless you have a contraindication for some other reason. However, you will need special counseling about the vaccine.		
Have you received monoclonal antibodies or convalescent plasma as part of COVID-19 treatment? Pfizer-BioNTech, Moderna, or Janssen COVID-19 vaccine should be deferred for at least 90 days to avoid interference of treatment with vaccine-induced immune responses.		
<18 years old: Have you had a well child visit with your pediatrician in the last 12 months?		

**NOTE:** Recipients of Janssen COVID-19 vaccine should be instructed to seek immediate medical attention if they develop shortness of breath, chest pain, leg pain or swelling, persistent abdominal pain, neurological symptoms (including severe or persistent headaches or blurred vision), nausea, vomiting, petechiae or easy bleeding beyond the site of vaccination within 4 to 30 days of receipt of Janssen vaccine. Most people who have developed blood clots and low platelets were females ages 18 through 49 years.

**NOTE:** Depending on vaccine type, a second dose of COVID-19 vaccine may be due in 21 days or 28 days after initial vaccine. Refer to your COVID-19 vaccination record card for second dose due date. Contact your vaccination provider, PCP, or your ADH Local Health Unit in 21 days or 28 days for more information. Keep your COVID-19 vaccination record card for your records for proof of initial vaccine date. Janssen COVID-19 vaccine is a ONE dose series.

**RELEASE AND ASSIGNMENT:** Please read the section on the reverse side of this form. The Providers Privacy Notice is available at the clinic site or accompanies this form. My signature indicates I have read, understand and agree to section **Release and Assignment** of the COVID-19 Immunization Consent Form and Vaccine Recipient Emergency Use of Authorization Fact Sheet (EUA).

**Signature of Patient/Parent/Guardian:**

\_\_\_\_\_

Date \_\_\_\_\_

